

Diagnostic Exercise

From The Davis-Thompson Foundation*

Case #: **163** Month: **April** Year: **2021**

Title: *Systemic Toxoplasmosis in a Red Kangaroo (Macropus rufus)*

Contributors: J. Pablo Velasco Montes de Oca,¹ DVM, Dipl. Cytopathology, MS student (Anatomic Pathology); Mariano Carossino,² DVM, PhD, DACVM; Ingeborg M. Langohr,² DVM, MS, PhD, DACVP; Udeni B. R. Balasuriya,² BVSc, MS, PhD, FSLCVS; Rudy Bauer,² DVM, PhD, DACVP

¹Departamento de Patología, Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, Ciudad de México, México.

²Louisiana Animal Diagnostic Disease Laboratory (LADDL) and Department of Pathobiological Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, LA, USA.




Figure 1: (A) Lateral view of the open thoracic cavity. (B) Lungs, dorsal view after being removed from the thorax.




Figure 2: (A) Lung. H&E, 50X. (B) Myocardium. H&E, 400X. Adapted from Carossino M, Bauer R, Mitchell MA, et al. Pathologic and immunohistochemical findings in an outbreak of systemic toxoplasmosis in a mob of red kangaroos. *Journal of Veterinary Diagnostic Investigation*. March 2021. doi:10.1177/10406387211001869




Figure 3: Lung. Alveolar spaces are diffusely filled by abundant histiocytes and alveolar edema. Pulmonary vessels and airways are delimited by cuffs composed of moderate numbers of lymphocytes, histiocytes and fewer plasma cells. H&E, 50X. Adapted from Carossino M, Bauer R, Mitchell MA, et al. Pathologic and immunohistochemical findings in an outbreak of systemic toxoplasmosis in a mob of red kangaroos. *Journal of Veterinary Diagnostic Investigation*. March 2021. doi:10.1177/10406387211001869




Figure 4: (A) Myocardium. Multifocal areas of the myocardium have fragmented cardiomyocytes, karyorrhectic debris, and mixed inflammatory cells including histiocytes, lymphocytes and plasma cells. H&E, 200X. (B) Myocardium. Occasionally within affected areas, cardiomyocytes contain intrasarcoplasmic protozoal cysts (arrows). H&E, 400X. Adapted from Carossino M, Bauer R, Mitchell MA, et al. Pathologic and immunohistochemical findings in an outbreak of systemic toxoplasmosis in a mob of red kangaroos. *Journal of Veterinary Diagnostic Investigation*. March 2021. doi:10.1177/10406387211001869

Histologic Description:

Lungs: The pulmonary parenchyma is diffusely and markedly congested, with alveolar spaces diffusely filled by abundant histiocytes and edema fluid. The pulmonary interstitium, most predominantly that around pulmonary vessels and occasional airways, is diffusely expanded by moderate numbers of lymphocytes, histiocytes and fewer plasma cells.

Heart: Multifocal areas throughout the myocardium are affected by necrosis characterized by foci of fragmented myofibers with increased sarcoplasmic eosinophilia and replacement by few histiocytes and lymphocytes, karyorrhectic debris, and proliferating fibroblasts. Occasional affected areas contain rare

intracellular protozoal cysts measuring approximately 15 x 5 µm and filled with numerous basophilic zoites.

Morphologic Diagnoses:

Lung: Pneumonia, interstitial, histiocytic, lymphocytic and plasmacytic, diffuse, moderate, subacute, with marked alveolar histiocytosis.

Heart: Myocarditis, necrotizing, multifocal, moderate, subacute, with intralumenal protozoal cysts.

Associated lesions/other organs affected:

Necrotizing myositis (skeletal muscle), necrotizing hepatitis, necrotizing encephalitis, lymphohistiocytic to necrotizing chorioretinitis and optic neuritis.

Possible causes:

- *Toxoplasma gondii*
- *Neospora caninum*
- *Trypanosoma cruzi* (depending on the geographic location)

Discussion:

Toxoplasma gondii is a zoonotic apicomplexan protozoal organism closely related to *Neospora caninum* and *Sarcocystis* spp., all of which are known to cause disease in numerous mammalian species. While all of these have similar life cycles, they involve different definitive hosts. For *T. gondii*, the definitive hosts are cats and other wild felids, in which the parasite completes an intestinal cycle leading to fecal shedding of infective oocysts into the environment. A diverse number of domestic and wild mammals can act as intermediary hosts for *T. gondii* (including but not limited to carnivores, ruminants, cetaceans, fishes, rodents, primates, and marsupials), which acquire the infection by ingestion of sporulated oocysts from the environment and can either remain subclinical or develop severe disease. The disease pattern is mostly associated with individual susceptibility as well as other host factors such as the immune response. Rapidly dividing tachyzoites disseminate systemically within the intermediate host, gradually undergoing encystation within tissues and switching to a slowly dividing cell phenotype (bradyzoites). Macropodids, including kangaroos, are known to be intermediate hosts highly susceptible to develop severe and systemic disease, like the case presented here. Infections have been reported in red (*Macropus rufus*), eastern grey (*Macropus giganteus*) and western grey kangaroos (*Macropus fuliginosus*). While a wide range of tissues can be affected, the muscular and nervous tissues are common sites where tissue cysts containing bradyzoites develop. Histologic and immunohistochemical differentiation of protozoa from the phylum Apicomplexa is

challenging due to the lack of distinct histomorphological features and extensive cross-reactivity among antibodies available for immunohistochemistry and immunofluorescence. Consequently, molecular methods are often necessary to reach a definitive diagnosis. The infection in kangaroos (as well as other intermediate hosts including humans) is acquired from the environment (food, water sources or soil) contaminated with infective cat feces. Therefore, the only means to control this disease in marsupial populations is to avoid contamination of the food/water supplies and environment by limiting access by outdoor cats or other wild felids into the premises. In humans, consumption of raw and undercooked meat as well as unpasteurized goat milk are important additional transmission routes of *T. gondii* and, consequently, mitigation strategies also focus on measures to guarantee food safety (proper cooking of meat products and consumption of pasteurized milk-derived products).

Trypanosoma cruzi (the agent associated with Chagas disease) is an important rule-out depending on the geographical location. This zoonotic protozoal organism is a non-apicomplexan, and its transmission cycle involves an insect vector (triatomine bug). Despite limited occurrence within the US, cases of *T. cruzi* infection are sporadically seen in companion animals (dogs), particularly in the Southern US. Histologically, a distinctive feature of *T. cruzi* amastigotes, which are preferentially found in the cardiac and skeletal muscles, is the presence of a kinetoplast adjacent to and parallel to its nucleus.

****This case is part of the following published study: Carossino M, Bauer R, Mitchell MA, et al. Pathologic and immunohistochemical findings in an outbreak of systemic toxoplasmosis in a mob of red kangaroos. *J Vet Diag Invest*. March 2021. doi:10.1177/10406387211001869**

References and Recommended Literature:

1. Carossino M, Bauer R, Mitchell MA, et al. Pathologic and immunohistochemical findings in an outbreak of systemic toxoplasmosis in a mob of red kangaroos. *J Vet Diag Invest*. March 2021. doi:10.1177/10406387211001869
2. Díaz-Ayala N, Hidalgo-Hermoso E, Cabello-Araya C, Carvallo-Chaigneau F. Infection with *Toxoplasma gondii* in a red kangaroo (*Macropus rufus*) and a Patagonian mara (*Dolichotis patagonum*) in captivity. *Rev Bras Parasitol Vet*. 2016;25(4):523-526. doi:10.1590/S1984-29612016076.
3. Hussain MA, Stitt V, Szabo EA, Nelan B. *Toxoplasma gondii* in the Food Supply. *Pathogens*. 2017;6(2):21. doi:10.3390/pathogens6020021
4. Kaufmann H, Yamage M, Roditi I, et al. Discrimination of *Neospora caninum* from *Toxoplasma gondii* and other apicomplexan parasites by hybridization and PCR. *Mol Cell Probes*. 1996;10(4):289-297. doi:10.1006/mcpr.1996.0038
5. Kim K, Weiss LM. *Toxoplasma gondii*: the model apicomplexan. *Int J Parasitol*. 2004;34(3):423-432. doi:10.1016/j.ijpara.2003.12.009

6. Miller DS, Faulkner C, Patton S. Detection of *Toxoplasma gondii* IgG antibodies in juvenile great grey kangaroos, *Macropus giganteus giganteus*. *J Zoo Wildl Med.* 2003;34(2):189-193. doi:10.1638/1042-7260(2003)034[0189:DOTGIA]2.0.CO;2
7. Müller N, Zimmermann V, Hentrich B, Gottstein B. Diagnosis of *Neospora caninum* and *Toxoplasma gondii* infection by PCR and DNA hybridization immunoassay. *J Clin Microbiol.* 1996;34(11):2850-2852.
8. Parameswaran N, O'Handley RM, Grigg ME, Fenwick SG, Thompson RC. Seroprevalence of *Toxoplasma gondii* in wild kangaroos using an ELISA. *Parasitol Int.* 2009;58(2):161-165. doi:10.1016/j.parint.2009.01.008
9. Su R, Dong H, Li T, et al. *Toxoplasma gondii* in four captive kangaroos (*Macropus* spp.) in China: Isolation of a strain of a new genotype from an eastern grey kangaroo (*Macropus giganteus*). *Int J Parasitol Parasites Wildl.* 2019;8:234-239. doi:10.1016/j.ijppaw.2019.03.003
10. Uzal F A, Plattner B, Hostetter, J M. Alimentary system. In: Maxie MG, ed. *Jubb, Kennedy and Palmers Pathology of Domestic Animals*. 6th ed. Vol. 2. Philadelphia, PA: Elsevier Saunders; 2016:235-236.

*The Diagnostic Exercises are an initiative of the **Latin Comparative Pathology Group (LCPG)**, the Latin American subdivision of The Davis-Thompson Foundation. These exercises are contributed by members and non-members from any country of residence. Consider submitting an exercise! A final document containing this material with answers and a brief discussion will be posted on the CL Davis website (http://www.cldavis.org/diagnostic_exercises.html).

Associate Editor for this Diagnostic Exercise: Mariano Carossino
Editor-in-chief: Claudio Barros